



King's Research Portal

DOI:

[10.1016/j.clnu.2017.08.028](https://doi.org/10.1016/j.clnu.2017.08.028)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Veronese, N., Stubbs, B., Noale, M., Solmi, M., Rizzoli, R., Vaona, A., Demurtas, J., Crepaldi, G., & Maggi, S. (2017). Adherence To A Mediterranean Diet Is Associated With Lower Incidence Of Frailty: A Longitudinal Cohort Study. *CLINICAL NUTRITION*. <https://doi.org/10.1016/j.clnu.2017.08.028>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Adherence To A Mediterranean Diet Is Associated With Lower Incidence Of Frailty: A Longitudinal Cohort Study

Nicola Veronese, Brendon Stubbs, Marianna Noale, Marco Solmi, Renè Rizzoli, Alberto Vaona, Jacopo Demurtas, Gaetano Crepaldi, Stefania Maggi



PII: S0261-5614(17)30311-4

DOI: [10.1016/j.clnu.2017.08.028](https://doi.org/10.1016/j.clnu.2017.08.028)

Reference: YCLNU 3240

To appear in: *Clinical Nutrition*

Received Date: 30 January 2017

Revised Date: 17 July 2017

Accepted Date: 30 August 2017

Please cite this article as: Veronese N, Stubbs B, Noale M, Solmi M, Rizzoli R, Vaona A, Demurtas J, Crepaldi G, Maggi S, Adherence To A Mediterranean Diet Is Associated With Lower Incidence Of Frailty: A Longitudinal Cohort Study, *Clinical Nutrition* (2017), doi: 10.1016/j.clnu.2017.08.028.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**ADHERENCE TO A MEDITERRANEAN DIET IS ASSOCIATED
WITH LOWER INCIDENCE OF FRAILTY:
A LONGITUDINAL COHORT STUDY**

Nicola Veronese^{1,2}, Brendon Stubbs^{3,4,5}, Marianna Noale¹, Marco Solmi^{2,6,7}, Renè Rizzoli⁸, Alberto
Vaona⁹, Jacopo Demurtas¹⁰, Gaetano Crepaldi¹, Stefania Maggi¹

¹ National Research Council, Neuroscience Institute, Aging Branch, Padova, Italy.

² Laboratory of Nutritional Biochemistry, Research Hospital, IRCCS “S. de Bellis”, Castellana
Grotte, Bari, Italy.

³ Physiotherapy Department, South London and Maudsley NHS Foundation Trust, Denmark Hill,
London SE5 8AZ, United Kingdom.

⁴ Health Service and Population Research Department, Institute of Psychiatry, Psychology and
Neuroscience King's College London, De Crespigny Park, London Box SE5 8AF, United Kingdom.

⁵ Faculty of Health, Social Care and Education, Anglia Ruskin University, Bishop Hall Lane,
Chelmsford CM1 1SQ, UK

⁶ Department of Neurosciences, University of Padova, Padova, Italy.

⁷ National Health Care System, Padova Local Unit ULSS 17, Italy.

⁸ Division of Bone Diseases, Department of Internal Medicine Specialties, Geneva University
Hospitals and Faculty of Medicine, Geneva, Switzerland.

⁹ Primary Care Department, Azienda ULSS20 Verona, Verona, Italy.

¹⁰ Primary Care Department, Azienda USL Toscana Sud Est, Grosseto, Italy.

23 **Address for correspondence:**

24 Nicola Veronese, MD

25 National Research Council, Neuroscience Institute, Aging Branch, Padova, Italy

26 Via Giustiniani, 2 35128 Padova, Italy

27 Phone: +390498218492; Fax: +390498211218

28 Email: ilmannato@gmail.com

ABSTRACT

Background & aims: There is a paucity of data investigating the relationship between the mediterranean and frailty, with no data among North American people. We aimed to investigate if adherence to a Mediterranean diet is associated with a lower incidence of frailty in a large cohort of North American people.

Methods: This study included subjects at higher risk or having knee osteoarthritis. Adherence to the Mediterranean diet was evaluated using a validated Mediterranean diet score (aMED) as proposed by Panagiotakos and classified into five categories. Frailty was defined using the Study of Osteoporotic Fracture (SOF) index as the presence of ≥ 2 out of: (i) weight loss $\geq 5\%$ between baseline and the subsequent follow-up visit; (ii) inability to do five chair stands; (iii) low energy level.

Results: During the 8 years follow-up, of the 4,421 participants initially included (mean age: 61.2 years, % of females=58.0), the incidence of frailty was approximately half in those with a higher adherence to the Mediterranean diet (8 for 1,000 person years) vs. those with a lower adherence (15 for 1,000 persons-years). After adjusting for 10 potential confounders (age, sex, race, body mass index, education, smoking habits, yearly income, physical activity level, Charlson co-morbidity index and daily energy intake), participants with the highest aMED scores were found to have a significant reduction in incident frailty (hazard ratio=0.71; 95% CIs: 0.50-0.99, $p=0.047$) with respect to those in a lower category. Regarding individual components of the Mediterranean diet, low consumption of poultry was found to be associated with higher risk of frailty.

Conclusions: A higher adherence to a Mediterranean diet was associated with a lower incidence of frailty over an 8-year follow-up period, even after adjusting for potential confounders.

Key words : frailty; Mediterranean diet; Osteoarthritis Initiative, older people

INTRODUCTION

Defined as “a state of increased vulnerability to stressors resulting from a decrease in physiologic reserves in multiple organ systems causing limited capacity to maintain homeostasis”,¹ frailty is a common condition in older people. It has been associated with an increased risk of several deleterious outcomes in that population, including disability, hospitalization and institutionalization.² Recent studies have also suggested that frailty could be considered an independent risk factor for cardiovascular^{3,4} and metabolic⁵ diseases that could further hasten the typical transition from frailty to disability.

Although the risk factors for frailty are numerous, diet could play its own relevant role in its development. Some observational studies have suggested that a deficiency of single nutrients, such as carotenoids, vitamins D, E, and C, folates, proteins, and whole grains^{6–12} might play a putative role in the development of frailty. But as the deficiency of some nutrients is often associated with the paucity of others, assessment of whole dietary patterns seems more appropriate for epidemiological purposes.¹³ Interestingly, a higher consumption of fruits and vegetables was associated with a reduced risk of frailty in a large cohort of older European subjects.¹⁴ The Mediterranean diet is one of the most important dietary patterns that has been investigated in view of its beneficial effects on several diseases. The term ‘Mediterranean diet,’ which describes traditional dietary habits of peoples from Mediterranean countries, is frequently depicted as a food pyramid¹⁵. For epidemiological purposes, adherence to the Mediterranean diet is usually evaluated using standardized questionnaires that take into consideration foods that are commonly eaten by Mediterranean populations. The Mediterranean-style diet is an established healthy-eating behavior that has consistently been shown to have beneficial effects on quality of life¹⁶ and on musculoskeletal¹⁷, cardiovascular¹⁸, metabolic¹⁹, and cognitive^{20,21} diseases.

79 Until now, to the best of our knowledge, only four studies have investigated the effect of the
80 Mediterranean diet on incident frailty.²²⁻²⁵ While all of these advanced the field and suggested that
81 the Mediterranean diet may play a role in delaying the onset of frailty, three were nevertheless
82 limited because of their short follow-up period^{22,23,25} (less than 4 years). No study has, moreover,
83 investigated the possible role of higher adherence to Mediterranean diet on frailty in a North
84 American population a population in which the higher presence of some medical conditions (such
85 as obesity)²⁶ that predicts frailty, could alter the association between Mediterranean diet and frailty.

86
87 Given the potential benefits of the Mediterranean diet on several diseases and the absence of data on
88 North American subjects, the current study aimed to investigate whether adherence to a
89 Mediterranean diet is associated with a lower incidence of frailty in a large cohort of North
90 Americans participating in the Osteoarthritis Initiative. We hypothesized that higher adherence to
91 the Mediterranean diet is associated with a lower incidence of frailty.

METHODS

Data source and subjects

Data were gathered from the Osteoarthritis Initiative (OAI)²⁷, available for public access at <http://www.oai.ucsf.edu/>. The specific datasets utilized for this research were registered during the baseline and screening evaluations (V00) and each database reporting data on frailty until 96 months from baseline (V10). Patients at high risk or having knee OA were recruited at four clinical centers in the USA (Baltimore, MD; Pittsburgh, PA; Pawtucket, RI; and Columbus, OH) between February 2004 and May 2006.

All the participants provided written informed consent. The OAI study protocol was approved by the institutional review board of the OAI Coordinating Center, University of California at San Francisco.

Adherence to Mediterranean diet (exposure)

Participants' diet patterns were analyzed using the Block Brief 2000 food frequency (FFQ) questionnaire during the baseline appointment.²⁸ The validated tool, containing a food list of 70 items, was designed to assess the individual's food and beverage consumption over the past year. Frequency of food consumption of the items included was reported at nine levels of intake from "never" to "every day". There were also seven dietary behavior questions on food preparation methods and fat intake, one question on fiber intake, and 13 questions on vitamin and mineral intakes.

Adherence to a Mediterranean diet was evaluated using the Mediterranean diet score (aMED) proposed by Panagiotakos et al.²⁹ The score was calculated based on a food frequency questionnaire completed during the baseline OAI visit. The aMED takes into consideration foods commonly consumed by individuals living in Mediterranean area. Participants were asked to rate their consumption of each single food item using a score ranging from 0 (less adherence) to 5 (better

adherence); the total possible score ranges from 0 to 55, with higher values indicating higher adherence to a Mediterranean diet. Cereals (e.g. bread, pasta, rice), potatoes, fruits, vegetables, legumes (e.g. peas, beans), fish were categorized on the basis of servings/month and specifically as: 0=never; 1=1 to 4 servings for month; 2=5 to 8; 3=9 to 12; 4=13 to 18; 5= more than 18 servings/month. As no distinction was made with reference to whole vs. refined cereals, all types of grains were considered under the same heading. Consumption of red meat, poultry, and full fat dairy products (e.g. milk cheese, yogurt) were categorized as: 0=more than 18 servings/month; 1=13 to 17 servings/month; 2=9 to 12; 3=5 to 8; 4=1 to 4; 5= never). Consumption of olive oil was categorized as the times it was used in a week's time and specifically as: 0=never; 1=rare; 2 \leq 1/weekly; 3= 2 times/weekly; 4=3 to 6; 5=daily. Alcoholic beverages were categorized as: 0 \geq 700 ml/day or 0; 1=600 to 699 ml/day; 2=500 to 599 ml/day; 3=400 to 499 ml/day; 4=300 to 399 ml/day; 5= $<$ 300 ml/day.

We arbitrary divided the population into 5 categories depending on their total scores as follows: aMED score <24 = Q1, 25-27 = Q2, 28-30 = Q3, 31-32 = Q4 and > 32 = Q5.

Outcome

The study's outcome of interest was incident frailty. In agreement with the Study of Osteoporotic Fracture (SOF) index^{30,31} frailty was defined as the presence of at least ≥ 2 out of three of the following criteria: (i) weight loss $\geq 5\%$ taking place between baseline and the follow-up examinations (at the baseline examination a body mass index, BMI, of less than 20 Kg/m², a common cut-off for identifying underweight people the elderly^{32,33}, was used, since no information regarding weight changes were recorded). Weight and height were measured at baseline and during follow-up examinations by a trained nurse; (ii) the inability to rise from a chair five times without arm support (hereafter referred to as inability to carry out chair stands); and (iii) poor energy based

on the SF12 questionnaire response of “little at a time” or “none at a time” to the question “in the past 4 weeks, did you have a lot of energy?”

The assessment of the outcome was made at the baseline and during the V01 (12 months), V03 (24 months), V05 (36 months), V06 (48 months), V08 (72 months) until the V10 (96 months).

Covariates

We identified 10 potential self-reported confounders including BMI; physical activity evaluated using the Physical Activity Scale for the Elderly (PASE);³⁴ race; smoking habit, educational level and yearly income ($<$ or \geq \$50,000 and missing data) to assess the relationship between aMED and incident frailty.

Validated general health measures of self-reported comorbidities were assessed using the modified Charlson comorbidity score³⁵. The medical morbidities that were assessed using that score regarded diseases/disorders that were common in North Americans such as fractures, heart attack and failure, stroke, chronic obstructive pulmonary disease, diabetes and cancer.³⁶

Statistical analyses

Normal distributions of quantitative variables were tested using the Kolmogorov-Smirnov test. Data are reported as means \pm standard deviations (SD) for continuous measures, and frequency and percentages for all discrete variables. Levene’s test was used to test the homoscedasticity of variances and, if its assumption was violated, Welch’s ANOVA was used. P values for trends were calculated using the Jonckheere-Terpstra test for continuous variables and the Mantel-Haenszel Chi-square test for categorical ones.

Cox’s regression analysis was used to explore the association between aMED at baseline and incident frailty. Deaths were censored and in a sensitivity analysis removed from the main analysis.

Factors significantly different across aMED categories (considering a p-value<0.10) or significantly associated with incident frailty at univariate analysis (p-value<0.05) were included. Multicollinearity among covariates was assessed using the variance inflation factor (VIF), with a score of 2 leading to the exclusion of a variable, but no parameter was excluded for this reason. The basic model was not adjusted for any confounders. The fully adjusted model included: age (as a continuous variable); sex; race (white vs. others); BMI (as continuous); education (degree vs. others); smoking habits (current and previous vs. others); yearly income (categorized as \geq or $<$ \$50,000 and missing data); Charlson comorbidity index; use of analgesic drugs (yes vs. no) and total energy intake (as a continuous variable). Cox's regression analysis data were reported as hazard ratios (HRs) with 95% confidence intervals (CIs).

We also ran the same analyses taking each single component of the Mediterranean diet as an outcome and classifying adherence as a low score (a score of 0, 1, or 2) or a high one (4-5). Adjusted HRs and 95% CI were in the end calculated to estimate the strength of the associations between aMED and incident frailty.

To test the robustness of our analyses, sensitivity analyses were conducted evaluating the interaction between aMED and selected factor (e.g. age below or more than 60 years, gender, race, education, smoking habits, yearly income and presence/absence of diseases at baseline) in predicting incident frailty, but no moderator emerged as significant.

All the analyses were performed using the SPSS 21.0 for Windows (SPSS Inc., Chicago, Illinois).

All statistical tests were two-tailed and statistical significance was assumed for a p-value <0.05 .

RESULTS

Sample selection

The OAI dataset includes a total of 4,796 North American participants. At baseline, 229 participants were excluded because data about their aMED adherence was missing or because they had implausible caloric intake (<500 or >5000 Kcal/day). Seventeen, who were already frail at baseline, were excluded and 129 were lost during the follow-up, leaving 4,421 participants whose data were included in the analyses. The excluded participants did not differ in terms of mean age and percentage of females compared to those included in this research.

Descriptive characteristics

Of the 4,421 participants, 1,857 were males and 2,564 females. Mean age was 61.2 years (± 9.2 years; range: 45-79). Mean aMED score was 28.1 points (± 5.1 points; range: 5-44).

Table 1 outlines the participants' characteristics classified into aMED categories. Those in the highest group (reflecting higher adherence to Mediterranean diet) were older (p for trend < 0.0001), more likely to be female (p for trend < 0.0001), white (p for trend < 0.0001), with a higher educational level (p for trend < 0.0001) with respect to those in the other categories. They also had lower BMI values (p for trend < 0.0001) and a lower prevalence of diabetes (p for trend < 0.0001) than other participants (**Table 1**).

Finally, those in the highest group, reported a significant lower presence of poor energy compared with the other participants (p for trend < 0.0001) (**Table 1**).

212 Table 1. Characteristics of the participants classified according to their adherence to a Mediterranean diet pattern.

	Q1 (n=1063) aMED<24	Q2 (n=884) aMED 25-27	Q3 (n=996) aMED 28-30	Q4 (n=597) aMED 31-32	Q5 (n=881) aMED > 32	P value for trend ^a
Energy intake (Kcal/day)	1407.1 (606.5)	1391.4 (557.7)	1437.8 (593.4)	1414.3 (541.7)	1417.7 (519.1)	0.31
Age (years)	58.9 (8.6)	61.1 (9.0)	61.6 (9.3)	62.7 (9.2)	63.0 (9.1)	<0.0001
PASE (points)	159.1 (84.1)	161.8 (79.9)	166.3 (83.9)	152.3 (77.6)	162.3 (82.0)	0.88
Females (n, %)	542 (51.0)	525 (59.4)	599 (60.1)	351 (58.8)	547 (62.1)	<0.0001
White race (n, %)	751 (70.7)	694 (78.5)	818 (82.2)	519 (86.9)	771 (87.6)	<0.0001
Smoking (previous/current) (n, %)	485 (45.9)	412 (46.8)	456 (46.1)	293 (49.2)	431 (49.2)	0.10
Graduate degree (n, %)	243 (22.9)	261 (29.6)	297 (29.8)	211 (35.3)	334 (37.9)	<0.0001
Yearly income (≥ \$50,000)	560 (52.7)	518 (58.6)	596 (59.8)	376 (63.0)	567 (64.4)	<0.0001
<i>Medical conditions</i>						
BMI (Kg/m ²)	29.8 (4.9)	29.2 (4.7)	28.4 (4.7)	28.1 (4.6)	27.4 (4.6)	<0.0001
Fractures (n, %)	166 (15.7)	166 (18.8)	165 (16.6)	124 (20.9)	165 (18.8)	0.12
Heart attack (n, %)	29 (2.8)	17 (2.0)	13 (1.3)	4 (0.7)	24 (2.8)	0.43

	Q1 (n=1063) aMED<24	Q2 (n=884) aMED 25-27	Q3 (n=996) aMED 28-30	Q4 (n=597) aMED 31-32	Q5 (n=881) aMED > 32	P value for trend ^a
Heart failure (n, %)	27 (2.6)	21 (2.4)	15 (1.5)	6 (1.0)	17 (1.9)	0.09
Stroke (n, %)	33 (3.1)	35 (4.0)	23 (2.3)	14 (2.4)	27 (3.1)	0.41
COPD (n, %)	27 (2.6)	19 (2.2)	23 (2.3)	9 (1.5)	18 (2.1)	0.30
Diabetes (n, %)	99 (9.6)	81 (9.3)	72 (7.3)	36 (6.1)	42 (4.8)	<0.0001
Cancer (n, %)	45 (4.2)	34 (3.8)	62 (6.2)	26 (4.4)	48 (5.5)	0.15
Frailty items at baseline						
Weight loss (n, %)	19 (1.8)	15 (1.7)	22 (2.2)	12 (2.0)	31 (3.5)	0.07
Inability to rise from a chair five times (n, %)	6 (0.6)	7 (0.8)	9 (0.9)	4 (0.7)	6 (0.7)	0.83
Poor energy (n, %)	160 (15.1)	110 (12.5)	100 (10.0)	53 (8.9)	67 (7.6)	<0.0001

213

214 **Notes:** The data are presented as means (with standard deviations) for continuous variables and number (with percentage).

215 ^a P values for trends were calculated using the Jonckheere-Terpstra test for continuous variables and the Mantel-Haenszel Chi-square test for

216 categorical ones.

217 **Abbreviations:** aMED: adherence to Mediterranean diet score; PASE: Physical Activity Scale for the Elderly; BMI: body mass index; COPD:
218 chronic obstructive pulmonary disease.

ACCEPTED MANUSCRIPT

Adherence to Mediterranean diet and incident frailty

During the 8 year follow-up, 362 (=8.2% of the baseline population) developed frailty for a global incidence rate of 13 (95%CI: 8-18)/1,000 persons-year.

Table 2 illustrates the association between aMED and frailty in the participants studied. The incidence of frailty was approximately half in those with higher aMED scores with respect to those with lower ones Q5: 8 (95%CI: 2-14) vs. Q1: 15, 95%CI: 8-22 for 1,000 persons-year, $p<0.0001$). Using Cox's regression analysis adjusting for 10 potential confounders, and taking those with the lowest adherence to Mediterranean diet as a reference (=Q1), the participants with the highest adherence to the Mediterranean diet had a significantly reduced probability of incident frailty of 29% (HR=0.71; 95% CI: 0.50-0.99, $p=0.047$; **Table 2**). However, the p for trend was not significant ($p=0.24$). Excluding 223 participants dead during the follow-up period attenuated our findings (adjusted HR for the highest quintile= 0.77; 95%CI: 0.54-1.10; $p=0.11$).

Of note, no moderator emerged as important in this association since the for interaction between aMED and age (more or below 60 years) was $p=0.17$, for gender $p=0.29$ and for presence of frailty criteria at baseline=0.17.

236 **Table 2. Association between adherence to Mediterranean diet and incidence of frailty.**

	Cumulative incidence (%)	Incidence (95% CI)	Unadjusted HR (95%CI)	P value	Fully-adjusted^a HR (95%CI)	P value
Q1 (aMED<24)	105/1063 (=9.9%)	15 (8-22)	1 [reference]		1 [reference]	
Q2 (aMED 25-27)	65/884 (=7.4%)	11 (4-18)	0.73 (0.54-1.00)	0.046	0.77 (0.56-1.05)	0.10
Q3 (aMED 28-30)	86/996 (=8.6%)	13 (6-20)	0.83 (0.63-1.11)	0.21	0.92 (0.68-1.23)	0.56
Q4 (aMED 31-32)	50/597 (=8.4%)	11 (3-19)	0.80 (0.57-1.12)	0.20	0.92 (0.65-1.30)	0.62
Q5 (aMED>32)	56/881 (=6.4%)	8 (2-14)	0.62 (0.45-0.85)	0.004	0.71 (0.50-0.99)	0.047

237 **Notes:**

238 All the data are presented as hazard ratios (HRs) with their 95% confidence intervals.

239 ^aFully-adjusted model included as covariates: age (as continuous); sex; race (whites vs. others); body mass index (as continuous); education (degree
240 vs. others); smoking habits (current and previous vs. others); yearly income (categorized as \geq or $<$ 50,000\$ and missing data); Physical Activity
241 Scale for Elderly score (as continuous); Charlson co-morbidity index; daily energy intake (in Kcal/die).

242 Abbreviations: CI: confidence intervals; HR: hazard ratio.

243 **Table 3** shows Cox's regression analyses of the single components of the Mediterranean diet. After adjusting for potential confounders, no
244 component was associated with a significantly lower incidence of frailty; lower consumption of poultry was associated with a higher risk of incident
245 frailty (HR=1.34, 95%CI: 1.07-1.67, p=0.0009).

246 **Table 3. Singular components of Mediterranean diet and incidence of frailty.**

	Number of events/number of participants (=cumulative incidence, %) [higher adherence; 4-5 points]	Number of events/number of participants (=cumulative incidence, %) [higher adherence; 0-3 points]	Unadjusted HR (95%CI)	P value	Fully-adjusted ^a HR (95%CI)	P value
Cereals	31/351 (8.8)	331/4070 (8.1)	0.89 (0.62-1.29)	0.55	0.80 (0.55-1.18)	0.26
Potatoes	311/3947 (7.9)	50/471 (10.6)	1.35 (1.00-1.82)	0.05	1.08 (0.79-1.48)	0.64
Fruits	107/1208 (8.9)	255/3212 (7.9)	0.87 (0.70-1.10)	0.24	0.86 (0.68-1.09)	0.20
Vegetables	24/186 (12.9)	337/4233 (8.0)	0.59 (0.38-0.89)	0.01	0.70 (0.45-1.08)	0.11
Legumes	290/3532 (8.2)	71/887 (8.0)	0.99 (0.77-1.29)	0.97	0.87 (0.66-1.14)	0.31
Fish	336/4020 (8.4)	25/375 (6.7)	0.82 (0.55-1.24)	0.35	0.77 (0.51-1.16)	0.21
Meat	320/3828 (8.4)	42/593 (7.1)	0.85 (0.62-1.17)	0.32	0.88 (0.63-1.23)	0.45

	Number of events/number of participants (=cumulative incidence, %) [higher adherence; 4-5 points]	Number of events/number of participants (=cumulative incidence, %) [higher adherence; 0-3 points]	Unadjusted HR (95%CI)	P value	Fully-adjusted ^a HR (95%CI)	P value
Poultry	187/2483 (7.5)	174/1912 (9.1)	1.21 (0.99-1.49)	0.07	1.34 (1.07-1.67)	0.009
Dairy	348/4196 (8.3)	14/224 (6.3)	0.81 (0.47-1.38)	0.44	0.64 (0.37-1.11)	0.12
Alcohol	224/2378 (9.4)	134/2027 (6.6)	0.68 (0.55-0.84)	<0.0001	0.85 (0.68-1.06)	0.15
Olive oil	311/3809 (8.2)	50/572 (8.7)	1.07 (0.79-1.44)	0.67	1.17 (0.86-1.58)	0.31

Notes:

All the data are presented as hazard ratios (HRs) with their 95% confidence intervals.

In all the analyses, we considered higher adherence to a component (as 4 or 5 points) vs. lower (0 to 3; reference).

^aFully-adjusted model included as covariates: age (as continuous); sex; race (whites vs. others); body mass index (as continuous); education (degree vs. others); smoking habits (current and previous vs. others); yearly income (categorized as \geq or $<$ 50,000\$ and missing data); Physical Activity Scale for Elderly score (as continuous); Charlson co-morbidity index; daily energy intake (in Kcal/die).

254 Abbreviations: CI: confidence intervals; HR: hazard ratio.

DISCUSSION

Our analysis of the OAI multi-center, longitudinal, prospective study of knee osteoarthritis in people living in North America uncovered that those who more closely followed a Mediterranean diet had a significantly lower incidence of frailty. After adjusting for 10 potential confounders, those with the highest aMED score had a significantly lower incidence of frailty by 29%. However, the significant association between Mediterranean diet and frailty disappears after excluding dead people, indicating that the Mediterranean diet is probably able to prevent severe forms of frailty and when considering the p for trends, suggesting that the association between aMED score and frailty is not necessarily linear.

Individuals reporting a higher adherence to a Mediterranean diet had significantly lower BMI values, a lower prevalence of diabetes and a higher education level with respect to the other participants. These factors could play a role in the lower incidence of frailty in individuals strictly following a Mediterranean diet. However, it should be noted that obese people tend to report significant lower calorie intake than those really introduced³⁷ and this may introduce a bias in our findings. There was also a significantly higher presence of two important risk factors for frailty in these participants, namely being female and older. This discrepancy, however, further confirmed a significant and independent association between higher adherence to this dietary pattern and lower incidence of frailty.

Several studies have examined the association between the intake of nutrients, and in particular protein intake,^{38,39} and the prevalence or incidence of frailty. A study including 1,345 older French participants reported that higher intake of proteins (and not of energy) was associated with a lower prevalence of frailty, also after taking in account several potential confounders.¹² A large study focusing on 24,417 women showed that increased protein intake is protective against the incidence of frailty.⁷ Other studies have reported, instead, that increased protein intake, particularly of an

animal origin, increases mortality rate^{40,41} or could be associated to an increased rate of oxidative stress in the intracellular compartment.^{42,43} Our study found conflicting results regarding the intake of animal proteins and incidence of frailty, since higher consumption of meat was not associated with the risk of frailty, although lower consumption of poultry did. Again, previous literature reported a clear association between higher proteins intake (independently from the source) and lower presence of frailty¹², suggesting that proteins are important in the prevention of frailty. The mechanism underlying this finding is unclear, but it is possible that poultry's nutritional characteristics (in particular its lower fat content compared to red meat) could play a role. Further research is needed to answer these questions.

Study results suggest that a Mediterranean diet is indeed associated with lower incidence of frailty, independently from potential confounders. The fact that a higher adherence to a Mediterranean diet was found to be linked to a decrease in inflammation⁴⁴ could mean that inflammation plays a role in the development of frailty⁴⁵ and that the anti-inflammatory properties linked to the phytochemicals present in foods contained in the diet are able to modify this association.⁴⁶ This hypothesis has been confirmed by other studies showing that higher use of antioxidant nutrients leads to lower incidence of frailty.^{6,8,9} In addition, several studies have demonstrated that a higher adherence to Mediterranean diet improves some protein pathways that are involved in the development of frailty, such as insulin resistance⁴⁷ and endothelial dysfunction.⁴⁸ Finally, a Mediterranean diet may reduce the risk of several chronic diseases, such as cardiovascular diseases, which may also lead to frailty.

Although other studies investigating the effect of higher adherence to Mediterranean diet on the development of frailty produced significant results,²²⁻²⁴ it is important to remember that they were carried out in continents other than North America, where quite different dietary patterns are utilized. The generalizability of these findings regarding North Americans people is thus

questionable. Furthermore, only one study had a long follow-up (six years) as the time window for frailty to develop.²⁴ The association between the Mediterranean diet and frailty uncovered by our data was thus confirmed by data linked to a large sample and with a long follow-up. Study results thus clearly indicate that this diet pattern is an important measure to prevent frailty and the conditions consequent to it, such as disability. Another important point is that these studies used the criteria suggested by Fried et al.⁴⁹ for defining frailty modified compared to the original version and other tools for assessing the adherence to Mediterranean diet, such as that suggested by Trichopoulou et al. in the EPIC study.⁵⁰ However, how these different definitions could affect the results of our and these studies is hard to determine.

The study does have some limitations, the main one being that we used a slightly different definition of frailty at baseline with respect to the one used at the follow-up as far as weight loss was concerned. Using that definition, only 17 participants were considered frail at baseline. Unfortunately, no data regarding weight changes were available before the enrollment in the OAI. Another concern regarding weight loss is that the reasons of this were not investigated and it is possible that some participants lost weight intentionally over the 8-year follow-up for improving the symptoms related to OA. Moreover, participants of the OAI are at high risk of OA at baseline and this could introduce another bias in our results. We were, moreover, unable to assess the influence of bio-humoral markers (e.g. inflammation, insulin-resistance) on the association between Mediterranean diet and frailty. In addition, the medical conditions were simply self-reported and not adjudicated by trained physicians. Fourth, we used a slightly modified version of the well-known Mediterranean diet adherence score²⁹. Finally, as we did not consider changes in dietary habits between baseline and the follow-up, this too could have introduced another bias.

In conclusion, our data analysis indicated that a higher adherence to a Mediterranean diet is associated with lower incidence of frailty, even after adjusting for several important confounders.

333 As far as single components were concerned, less consumption of poultry was associated with
334 higher risk of frailty; this could suggest that animal origin proteins should be included in an
335 optimum Mediterranean diet. Future interventional studies are warranted to confirm our findings.

ACKNOWLEDGEMENTS

Conflict of interest: None of the authors have any financial arrangements, organizational affiliations or other relationships that might give rise to any conflict of interest regarding the subject matter of the manuscript submitted.

Founding source: The OAI is a public-private partnership comprised of five contracts (N01-AR-2-2258; N01-AR-2-2259; N01-AR-2-2260; N01-AR-2-2261; N01-AR-2-2262) funded by the National Institutes of Health, a branch of the Department of Health and Human Services, and conducted by the OAI Study Investigators. Private funding partners include Merck Research Laboratories; Novartis Pharmaceuticals Corporation, GlaxoSmithKline; and Pfizer, Inc. Private sector funding for the OAI is managed by the Foundation for the National Institutes of Health. This manuscript was prepared using an OAI public use data set and does not necessarily reflect the opinions or views of the OAI investigators, the NIH, or the private funding partners.

Sponsor's role: the sponsors had no role in designing the study, in patient recruitment, data collection/analysis or in drafting the manuscript.

Authors' contributions: Study concept and design: Veronese, Stubbs. Data analysis and interpretation: Noale, Veronese, Solmi. Drafting of the manuscript: Vaona, Demurtas, Veronese, Stubbs. Critical revision of the manuscript: Maggi, Rizzoli. Statistical analysis: Veronese, Noale.

Conflict of interest: none.

REFERENCES

1. Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K. Frailty in elderly people. *Lancet*. 2013;381(9868):752-762. doi:10.1016/S0140-6736(12)62167-9.
2. Fugate Woods N, LaCroix AZ, Gray SL, et al. Frailty: Emergence and Consequences in Women Aged 65 and Older in the Women's Health Initiative Observational Study. *J Am Geriatr Soc*. 2005;53(8):1321-1330. doi:10.1111/j.1532-5415.2005.53405.x.
3. Afilalo J, Alexander KP, Mack MJ, et al. Frailty assessment in the cardiovascular care of older adults. *J Am Coll Cardiol*. 2014;63(8):747-762. doi:10.1016/j.jacc.2013.09.070.
4. Sergi G, Veronese N, Fontana L, et al. Pre-frailty and risk of cardiovascular disease in elderly men and women: the pro.v.a. study. *J Am Coll Cardiol*. 2015;65(10):976-983. doi:10.1016/j.jacc.2014.12.040.
5. Veronese N, Stubbs B, Fontana L, et al. Frailty Is Associated with an Increased Risk of Incident Type 2 Diabetes in the Elderly. *J Am Med Dir Assoc*. June 2016. doi:10.1016/j.jamda.2016.04.021.
6. Sandoval-Insausti H, Pérez-Tasigchana RF, López-García E, García-Esquinas E, Rodríguez-Artalejo F, Guallar-Castillón P. Macronutrients Intake and Incident Frailty in Older Adults: A Prospective Cohort Study. *J Gerontol A Biol Sci Med Sci*. 2016;0(0):1-6. doi:10.1093/gerona/glw033.
7. Beasley JM, LaCroix AZ, Neuhaus ML, et al. Protein intake and incident frailty in the women's health initiative observational study. *J Am Geriatr Soc*. 2010;58(6):1063-1071. doi:10.1111/j.1532-5415.2010.02866.x.
8. Michelson E, Blaum C, Semba RD, Xue Q-L, Ricks MO, Fried LP. Vitamin and carotenoid status in older women: associations with the frailty syndrome. *J Gerontol A Biol Sci Med Sci*. 2006;61(6):600-607. doi:10.1111/j.1532-5415.2010.02866.x.
9. Semba RD, Bartali B, Zhou J, Blaum C, Ko CW, Fried LP. Low serum micronutrient concentrations predict frailty among older women living in the community. *Journals*

Gerontol - Ser A Biol Sci Med Sci. 2006;61(6):594-599.

10. Bartali B, Frongillo EA, Bandinelli S, et al. Low nutrient intake is an essential component of frailty in older persons. *Journals Gerontol - Ser A Biol Sci Med Sci.* 2006;61(6):589-593.

11. Wilhelm-Leen ER, Hall YN, Deboer IH, Chertow GM. Vitamin D deficiency and frailty in older Americans. *J Intern Med.* 2010;268(2):171-180. doi:10.1111/j.1365-2796.2010.02248.x.

12. Rahi B, Colombet Z, Gonzalez-Colaço Harmand M, et al. Higher Protein but Not Energy Intake Is Associated With a Lower Prevalence of Frailty Among Community-Dwelling Older Adults in the French Three-City Cohort. *J Am Med Dir Assoc.* 2016;17(7):672.e7-672.e11. doi:10.1016/j.jamda.2016.05.005.

13. Jacobs DR, Steffen LM. Nutrients, foods, and dietary patterns as exposures in research: A framework for food synergy. In: *American Journal of Clinical Nutrition.* Vol 78. ; 2003.

14. García-Esquinas E, Rahi B, Peres K, et al. Consumption of fruit and vegetables and risk of frailty: A dose-response analysis of 3 prospective cohorts of community-dwelling older adults. *Am J Clin Nutr.* 2016;104(1):132-142. doi:10.3945/ajcn.115.125781.

15. Willett WC, Sacks F, Trichopoulos A, et al. Mediterranean diet pyramid: A cultural model for healthy eating. *Am J Clin Nutr.* 1995;61(6 SUPPL.). doi:10.1007/s13398-014-0173-7.2.

16. Veronese N, Stubbs B, Noale M, Solmi M, Luchini C, Maggi S. Adherence to the Mediterranean diet is associated with better quality of life: data from the Osteoarthritis Initiative. *Am J Clin Nutr.* September 2016:1-6. doi:10.3945/ajcn.116.136390.

17. Haring B, Crandall CJ, Wu C, et al. Dietary Patterns and Fractures in Postmenopausal Women. *JAMA Intern Med.* March 2016. doi:10.1001/jamainternmed.2016.0482.

18. Nissensohn M, Román-Viñas B, Sánchez-Villegas A, Piscopo S, Serra-Majem L. The Effect of the Mediterranean Diet on Hypertension: A Systematic Review and Meta-Analysis. *J Nutr Educ Behav.* 2016;48(1):42-53.e1. doi:10.1016/j.jneb.2015.08.023.

19. Schwingshackl L, Missbach B, König J, Hoffmann G. Adherence to a Mediterranean diet and

risk of diabetes: a systematic review and meta-analysis. *Public Health Nutr.*

2015;18(7):1292-1299. doi:10.1017/S1368980014001542.

20. Cao L, Tan L, Wang H-F, et al. Dietary Patterns and Risk of Dementia: a Systematic Review and Meta-Analysis of Cohort Studies. *Mol Neurobiol.* November 2015. doi:10.1007/s12035-015-9516-4.

21. Lourida I, Soni M, Thompson-Coon J, et al. Mediterranean diet, cognitive function, and dementia: a systematic review. *Epidemiology.* 2013;24(4):479-489. doi:10.1097/EDE.0b013e3182944410.

22. León-Muñoz L, Guallar-Castillón P, López-García E, Rodríguez-Artalejo F. Mediterranean Diet and Risk of Frailty in Community-Dwelling Older Adults. *J Am Med Dir Assoc.* 2014;15(12):899-903. doi:10.1016/j.jamda.2014.06.013.

23. Chan R, Leung J, Woo J. Dietary patterns and risk of frailty in Chinese community-dwelling older people in Hong Kong: A prospective cohort study. *Nutrients.* 2015;7(8):7070-7084. doi:10.3390/nu7085326.

24. Talegawkar SA, Bandinelli S, Bandeen-Roche K, et al. A higher adherence to a Mediterranean-style diet is inversely associated with the development of frailty in community-dwelling elderly men and women. *J Nutr.* 2012;142(12):2161-2166. doi:10.3945/jn.112.165498.

25. Rahi B, Ajana S, Tabue-Teguo M, Dartigues J-F, Peres K, Fear C. High adherence to a Mediterranean diet and lower risk of frailty among French older adults community-dwellers: Results from the Three-City-Bordeaux Study. *Clin Nutr.* May 2017. doi:10.1016/j.clnu.2017.05.020.

26. Veronese N, Cereda E, Solmi M, et al. Inverse relationship between body mass index and mortality in older nursing home residents: a meta-analysis of 19,538 elderly subjects. *Obes Rev.* 2015;16(11):1001-1015. doi:10.1111/obr.12309.

27. Felson DT, Nevitt MC. Epidemiologic studies for osteoarthritis: New versus conventional

study design approaches. *Rheum Dis Clin North Am*. 2004;30(4):783-797.

doi:10.1016/j.rdc.2004.07.005.

28. Block G, Hartman AM, Naughton D. A reduced dietary questionnaire: development and validation. *Epidemiology*. 1990;1(1):58-64.

29. Panagiotakos DB, Pitsavos C, Stefanadis C. Dietary patterns: a Mediterranean diet score and its relation to clinical and biological markers of cardiovascular disease risk. *Nutr Metab Cardiovasc Dis*. 2006;16(8):559-568. doi:10.1016/j.numecd.2005.08.006.

30. Ensrud KE, Ewing SK, Cawthon PM, et al. A comparison of frailty indexes for the prediction of falls, disability, fractures, and mortality in older men. *J Am Geriatr Soc*. 2009;57(3):492-498. doi:10.1111/j.1532-5415.2009.02137.x.

31. Misra D, Felson DT, Silliman RA, et al. Knee osteoarthritis and frailty: findings from the Multicenter Osteoarthritis Study and Osteoarthritis Initiative. *J Gerontol A Biol Sci Med Sci*. 2015;70(3):339-344. doi:10.1093/gerona/glu102.

32. Veronese N, De Rui M, Toffanello ED, et al. Body Mass Index as a Predictor of All-Cause Mortality in Nursing Home Residents During a 5-Year Follow-up. *J Am Med Dir Assoc*. 2013;14(1):53-57. doi:10.1016/j.jamda.2012.09.014.

33. Diekmann R, Winning K, Uter W, et al. Screening for malnutrition among nursing home residents - a comparative analysis of the mini nutritional assessment, the nutritional risk screening, and the malnutrition universal screening tool. *J Nutr Health Aging*. 2013;17(4):326-331. doi:10.1007/s12603-012-0396-2.

34. Washburn RA, McAuley E, Katula J, Mihalko SL, Boileau RA. The physical activity scale for the elderly (PASE): evidence for validity. *J Clin Epidemiol*. 1999;52(7):643-651. <http://www.ncbi.nlm.nih.gov/pubmed/10391658>. Accessed March 17, 2016.

35. Katz JN, Chang LC, Sangha O, Fossel AH, Bates DW. Can comorbidity be measured by questionnaire rather than medical record review? *Med Care*. 1996;34(1):73-84. <http://www.ncbi.nlm.nih.gov/pubmed/8551813>. Accessed March 25, 2016.

- 462 36. Mahmood SS, Levy D, Vasan RS, Wang TJ. The Framingham Heart Study and the
 463 epidemiology of cardiovascular disease: a historical perspective. *Lancet*.
 464 2016;383(9921):999-1008. doi:10.1016/S0140-6736(13)61752-3.
- 465 37. Kye S, Kwon S-O, Lee S-Y, et al. Under-reporting of Energy Intake from 24-hour Dietary
 466 Recalls in the Korean National Health and Nutrition Examination Survey. *Osong Public Heal*
 467 *Res Perspect*. 2014;5(2):85-91. doi:10.1016/j.phrp.2014.02.002.
- 468 38. Artaza-Artabe I, Sáez-López P, Sánchez-Hernández N, Fernández-Gutierrez N, Malafarina
 469 V. The relationship between nutrition and frailty: Effects of protein intake, nutritional
 470 supplementation, vitamin D and exercise on muscle metabolism in the elderly. A systematic
 471 review. *Maturitas*. April 2016. doi:10.1016/j.maturitas.2016.04.009.
- 472 39. Deutz NEP, Bauer JM, Barazzoni R, et al. Protein intake and exercise for optimal muscle
 473 function with aging: Recommendations from the ESPEN Expert Group. *Clin Nutr*.
 474 2014;33:929-936. doi:10.1016/j.clnu.2014.04.007.
- 475 40. Ligiou P, Sandin S, Weiderpass E, et al. Low carbohydrate?high protein diet and mortality in
 476 a cohort of Swedish women. *J Intern Med*. 2007;261(4):366-374. doi:10.1111/j.1365-
 477 2796.2007.01774.x.
- 478 41. Levine ME, Suarez JA, Brandhorst S, et al. Low Protein Intake Is Associated with a Major
 479 Reduction in IGF-1, Cancer, and Overall Mortality in the 65 and Younger but Not Older
 480 Population. *Cell Metab*. 2014;19(3):407-417. doi:10.1016/j.cmet.2014.02.006.
- 481 42. Fontana L, Cummings NE, Arriola Apelo SI, et al. Decreased Consumption of Branched-
 482 Chain Amino Acids Improves Metabolic Health. *Cell Rep*. June 2016.
 483 doi:10.1016/j.celrep.2016.05.092.
- 484 43. Rizza W, Veronese N, Fontana L. What are the roles of calorie restriction and diet quality in
 485 promoting healthy longevity? *Ageing Res Rev*. 2014;13(1):38-45.
 486 doi:10.1016/j.arr.2013.11.002.
- 487 44. Chrysohoou C, Panagiotakos DB, Pitsavos C, Das UN, Stefanadis C. Adherence to the

- Mediterranean diet attenuates inflammation and coagulation process in healthy adults: The ATTICA Study. *J Am Coll Cardiol.* 2004;44(1):152-158. doi:10.1016/j.jacc.2004.03.039.
45. Soysal P, Stubbs B, Lucato P, et al. Inflammation And Frailty In The Elderly: A Systematic Review And Meta-analysis. *Ageing Res Rev.* 2016. doi:10.1016/j.arr.2016.08.006.
46. Musumeci G, Trovato FM, Pichler K, Weinberg AM, Loreto C, Castrogiovanni P. Extra-virgin olive oil diet and mild physical activity prevent cartilage degeneration in an osteoarthritis model: An in vivo and in vitro study on lubricin expression. *J Nutr Biochem.* 2013;24(12):2064-2075. doi:10.1016/j.jnutbio.2013.07.007.
47. Salas-Salvadó, Jordi, Bullo M BN. Reduction in the Incidence of Type 2 Diabetes With the Mediterranean Diet. *Diabetes Care.* 2011;34(1):14-19. doi:10.2337/dc10-1288.
48. Kok FJ, Kromhout D. Atherosclerosis--epidemiological studies on the health effects of a Mediterranean diet. *Eur J Nutr.* 2004;43 Suppl 1:I/2-5. doi:10.1007/s00394-004-1102-x.
49. Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci.* 2001;56(3):M146-56.
<http://www.ncbi.nlm.nih.gov/pubmed/11253156>. Accessed August 27, 2014.
50. Trichopoulou A, Orfanos P, Norat T, et al. Modified Mediterranean diet and survival: EPIC-elderly prospective cohort study. *BMJ.* 2005;330(7498):991.
<http://www.bmj.com/content/330/7498/991.abstract>.